1	"Rare Poultry Parasites." p. 1310 (ZA SOCIALISTICKE ZEMEDEISTVI, Vol. 3, No. 11, Nov. 1953) Praha, Czechoslovakia															
٠.						-										
	SO:	Month April	ly Lis 1954.	st of	East lassi	Europe:	an Ac	cession	15, Li	orary o	of Cong	ress,	Vol.	3, N	0. 4,	
•		Apr As	. 1//4													
				-												. •
							•	-								 

CZECHOSLOVAKIA/Diseases of Farm Animals - Diseases Caused by

R-3

Helminths.

: Ref Zhur - Biol., No 11, 1950, 50239 Abs Jour

Author

Willomitzer, J.

Inst

Title

The Diagnosis of Notocotyliasis in Ducks.

Orig Pub

Veterinarstvi, 1957, 7, No 9, 268-269

Abstract

: A report is given of a notocotyliasis infection in ducks (causative agent is Notocotylus attennatus). Out of 580 ducks 10 died. In the perished ducks, cachexia, enteritis, and degeneration of parenchymatous organs were noted. A morphological description of the parasite is given.

Card 1/1

CCUPTRY : CZECHOSLOVAKIA

CATECORY : Zooparasitology. Parasitic Protozoa. Flagellata

ABS. JOUR. : RZhBiol., No. 2 1959, No. 5677

AUTHOR : Willoritzeracy.

IIST. :

TITLE : Morphological Studies of Trichomonas suis

ORIG. PUB. : Veterinarstvi, 1958, 8, No 5, 167-168

ABSTRACT : No abstract

WILLOMITZER, J., dr., C.Sc.; LEBDUSKA, J., prof., dr., Dr.Sc.

Treatment of poultry capillariasis by Methyridin and other drugs. Vestnik vyzk zemedel 9 no.10:471-473 '62.

1. Vyzkumny ustav veterinarniho lekarstvi, Brno.

P

CZECHOSLOVAKIA

WILLOMITZER, J., DVM, Cand of Scien; LEBDUSKA, J., Prof. DVM, Dr. of Scien.

Brno (for both)

Prague, Veterinarstvi, No 6, 1963, pp 257-259

"Influence of Invasion of Capillaria caudinflata on the Weight of Hens."

WILLOMITZER, Jindrich, MVDr. CSc.

Effect of Czechoslovak anticestodiasis preparations on hen egg laying. Vest ust zemedel 11 no. 7:262-264 Jl '64.

1. Research Institute of Voterinary Medicine, Brno.

WILLOMITZER, Jindrich, MVDr. CSc.; LEBDUSKA, Jaroslav, prof. MVDr. DrSc.

Experiences in the treatment of poultry capillariosis by methyridine.
Veter medicina 8 no.4:245-248 '63.

1. Research Institute of Veterinary Medicine, Brno. Director of the
Institute [doc. MVDr. inz.] Jan Vlcek.

WILLOMITZER, Jindrich, MVDr. CSC.

Tests of a more suitable method of organic tir compound application in poultry infection by Davainea proglottina. Veter medicina 8 no.4: 249-256 163.

1. Research Institute of Veterinary Medicine, Brno. Director of the Institute [doc. MVDr. inz.] Jan Vlcek.

MELICHAR, Bohuslav, doc. dr. PhMr.; WILLOMITZER, Jindrich, MVDr. CSc.

Effect of aerosols of nontoxic physical insecticides on poultry ectoparasites. Veter medicina 8 no.4:275-284 163.

1. Research Institute of Veterinary Medicine, Brno. Director of the Institute [doc. MVDr. inz.] Jan Vlcek.

CZECHOSLOVAKIA

WILLOMITZER, Jindrich, MVDr, CSc

Brno

Brno, Veterinarstvi, No 3 [March] 1967, pp 116-117

"On therapy and chromoprophylaxis of some paratosis in scratching fowl."

MELLCHAR, Bohnslav, Docent, Dr. Hagdester of Pharmacy, and Willo-Mitaer, Jindrich, Dr of Veterinary Medicine, Candidate of Sciences, Research Institute of Veterinary Medicine (Vyzkumny ustav veterinarniho lekarstvi), Brno, Docent Engr Jan VLCEK, Dr of Veterinary Medicine, director.

"Effect of the Aerosols of Montoxic Physical Insecticides on the Ectoparasites of Fowl."

Prague, Veterinarni Hedicina, Vol &(XXXVI), Ho 4, August 63, pp 275-234.

Abstract [Authors' English summary, modified]: Determined were the physical and chemical parameters of ten substances of the following types: SiO<sub>2</sub> aerogels, silicates, adsorption carbon, and fly ashes. A method was established for determining the effect of physical insecticides on Dermanyssus gallinae and Musca dom. The correlation was determined between the size of the surface of the SiO<sub>2</sub> substances and silicates and their effect on Dermanyssus in vitro. The nontoxicity of three SiO<sub>2</sub> aerogels in chickens is reported as well as their positive effect on Dermanyssus gallinae and Menopon gallinae in hens. Futher experiments are announced on the evolution cycles and techniques of application. Twenty references, including 8 Czech and 1 Russian.

CAECHOSLOVAKIA

WILLOHITZER, Jindrich, Dr of Veterinary Medicine, Candidate of Sciences, Research Institute of Veterinary Medicine (Vyzkumny ustav veterinarniho lekarstvi), Brno, Docent Engr Jan VLCEK, Dr of Veterinary Medicine, director.

"Trials With a More Suitable Way of Applying the Organic Tin Compound in the Davaineosis of Poultry."

Frague, <u>Veterinarni Medicina</u>, Vol 8(XXXVI), No 4, August 63, pp 249-256.

Abstract [Author's English summary, modified]: The trials have corroborated a relatively good effectiveness of the stable Dinnbutyl tin dilaurate preparations made in Czechoslovakia. Data are given on experiments with Davainex 10 and Davainex 25. A comparison of the two preparations showed that an application of Davainex 10 in doses substantially lower than Davainex 25 produced only a slightly lower effect. Experiments are under way to determine the effect of Davainex 10 on the output of eggs of animals under treatment. Four references, including 1 Czech.

1/1


Sciences, and LEBDUSKA, Jaroslav, Prof., Dr of Veterinary Medicine, Candidate of dicine, Dr of Sciences, Research Institute of Veterinary Medicine (Vyzkumny ustav veterinarni mediciny)

### WIIMANSKA, A.; BASINSKI, A.

"Influence of Electrolytes on the Adsorption of Sulfurous Acid by Potato Starch." P. 82. (PRZEMISL CHEMICZNI, Vol. 10, No. 2, Feb. 1954, Warszawa, Poland)

SO: Monthly List of East European Accessions, (EEAL), LC, Vol. 4, No. 1, Jan. 1955 Uncl.

# WILMANSKI, Bernard Studies on the influence of the balloon control ring upon the stressing of the cotton yarn spun on cotton frames. Przegl wlokien 16 no.2:73-79 F '62. 1. Centralne Biuro Technicane Przemyslu Maszyn Wlokienniczych, Lodz.



SCHMANN, Jadwiga; WILMANSKA, Jadwiga; WINIARSKI, Waclaw

Erythroderma and other cutaneous complications in gold therapy.

Przegl. derm. 48 no.4:341-348 JI-14g. 61.

1. Z Kliniki Dermatolegicznej PAM w Szczecinie Kierownik: doc. dr

H. Prochackt. Z Kliniki Chorob Wedne trznych PAM w Szczecinie Kierownik:

pref. dr F. Belechowski.


(GOLD texicol) (EKYTHRODERMA etiol)

(DERMATITIS MEDICAMENTOSA)

WILMANSKA, Jadwiga

Studies on chemical properties of trichophytin. Roczn. pom. akad. med. Swierczewski 11:525-533 '165.

1. Z Kliniki Dermatologicznej Pomorskiej Akademii Medycznej (Kierownik: prof. dr. med. Henryk Prochacki).



44925 P/006/62/010/004/003/004 D237/D308

Wilmański Krzysztof

TITLE:

Elastic-plastic stresses in a ring, due to a temper-

ature field

PERIODICAL:

Rozprawy Inzynierskie, v. 10, no. 4, 1962, 713-729

The author discusses the elastic-plastic state of stress in a ring under a steady state radial temperature field. Saint-Venant model is assumed. Using Treska yield condition the author shows that one or two plastic regions may appear depending on the magnitude of the coefficient  $E\alpha_T$  |  $\Delta T$  |  $/\sigma_s$ , where E - Young's modulus;  $\alpha_T$  - coeff. of thermal expansion;  $\Delta T$  -  $T_a$  -  $T_b$ ; a and b - inner and outer radius of the ring; os - yield point. The inner plastic zone between r = a, r = 5 always appears first, irrespective of the sign of  $\Delta T$ . The outer plastic zone is bounded, when it appears, by  $r = \eta$ , r = 0. Expressions for radial and tangential stresses and the condition for the appears r = 0. tion for the appearance of the outer plastic zone in terms of 5 are

Card 1/2

Elastic-plastic stresses

P/006/62/010/004/003/004 D237/D308

found. Stresses in the ring in the presence of plastic zones and formulas for 5 and 7 are given in terms of Bessel functions and are shown to be reducible to the relations given by J.C. Wilhoit, Jr. The results obtained may be considered to be sufficiently accurate for temperatures up to 350°C, while for higher temperatures they may be regarded as first approximations to exact solutions. Comparison is made with the results of J.C. Wilhoit, Jr. There are 7 figures.


ASSOCIATION:

Politechnika Zódzka (Zódź Polytechnic)


SUBMITTED:


April 9, 1962


Card 2/2




"APPROVED FOR RELEASE: 03/20/2001 CIA-RDP86-00513R001961620001-5









POLAND / General Division, Problems of Teaching A-8

Abs Jour: Ref Zhur-Biologila, No 5, 1958, 18955

Author: Wilska Z.

Inst :

Title : The Plan of Work of a Circle of Amateur Fish Breeders

in Aquariums

Orig Pub: Biol. szkole, 1957, 10, No 2, 107-109

Abstract: No abstract

Card 1/1

### WILSKA-JESZKA, J.

The chemistry and physiological properties of flavonoids. p. 289.

WIADOMOSCI CHEMICZNE. (Polskie Towaezystwo Chemiczne) Wroclaw, Poland. Vol. 13, no. 6, June 1959.

Monthly List of East European Accessions (EEAI) LC, Vol. 9, no. 1, Jan. 1960.

Uncl.


POLAND


### WILSKA-JESZKA, Jadwiga

Department of Nutritives and Vitamin Concentrate Technology, Lodz Polytechnic (Katedra Technologii Odzywek i Koncentratow Witaminowych Politechniki Lodzkiej)

Wroclaw, Wiadomosci chemiczne, No 1, January 1966, pp 72-72

"Characteristics of flavonic compounds occurring in the Sorbus acuparia L. fruits." (doctoral thesis)





Vol. 6,	No. 3,	March 1953	. Warsza	ies in low wa.)						
		# #								
a file Having til som st Filesaal til anti-					. · · · · · · · · · · · · · · · · · · ·					
									. ·	
				en Accessio	na Vol	3. No. 2,	Library	of Congress	3,	

Zooparasitology. Parasitic Worms. Plant Country CATEGORY 1959, No. 10304 Helminths RZBiol., No. A3S. JOUR. Wilski Andrzej AUTHOR Research on the Control of the Beet Nematode by MSI. TIPLE Chemical Means Roczn. nauk rolniczych, 1957, A 75, No 4, 645-666 ORIG. PUB. : In small bowls and under field conditions experiments were peformed on the testing of ABSTRACT dinitroorthocresol (1) (41 and 50% of the active principle), intox (chlordane), dichlorethane, forbiate, chlorphenol, wolatox and systox for the purpose of disinfecting the soil and plants with respect to the best nematode. I was tested in doses of 7.2-80 kilograms/hecture; it was introduced into the soil in an aqueous solution 1 to 4 days before sceding the bects. The efficacy of I (reduction in the number of female 1/3 CARD:

COURTRY CATEGORY 1959, No. 10304 A85. JOUR. : RZB101., No. ROHTUA TROT. ITTI CRIG. PUB. : : nematodes on the roots) did not exceed 80%. Chlorphenol in a dose of 5 liters per square ABSTRACT centimeter in a 1 to 5% solution and 25 liters of a 1% solution did not give satisfactory results either. In addition, the use of solutions was practically impossible because of the large quantities of water required. Dichlorethane (5-20 tons/hectare) did not cause any reduction in the number of nematodes, but an increase in the beet yield was observed. Forbiate (0.5-2 tons/hectare) did not exert any toxic effect on CARD: 2/3

SAN DESCRIPTION OF THE STREET OF THE STREET

Country CATEGORY ABS. JOUR. : REBlol., No. 1959, No. 10304 AUTHOR INST. TITLE ORIG. PUB. : ABSTRACT the nematode under field conditions compared with the flower pots, which is explained by the poorer mixing of the preparation with the soil in the field. In doses of 600-1200 cubic centimeters/ hectare intox did not give any satisfactory results; the same thing was observed with wofatox (100-500 kilograms/hectare) and sprinkling with a 1% system solution. Of all the preparations tested forbiate has the least number of harmful properties. Bibliography 54 titles. --N. M. Sveshnikova CARD: 3/3

89185 P/005/60/000/052/002/003 A076/A026

26.2182

Wilski, Artur, Master of Engineering

TITLE:

AUTHOR:


Lubricating Technique Today


PERIODICAL:

Przeglad Techniczny, 1960, No. 52, p. 5

TEXT: The author describes in general properties of lubricants used in rockets, inter-continental mussiles, jet aircraft and engines operating at high temperature. The surface of a super-sonic aircraft during flight heats up to 130 - 530°C presenting a lubricating problem. New types of lubricant are needed. The most important lubricants are based on various types of silicons. Silicon lubricants have the best lubricating properties, e.g. low steam tension, resistance to high temperature, good viscosity, indifferent to changes in temperature, they do not corrode and may be used as shock and heat absorbing liquid. A higher class of lubricant used in aircraft are the so-called esters of aromatic acid. Practical application of phenylbicarboxyl acid esters gave good results, as this type of lubricant is not affected by hydrolysis, has low steam tension and does not yield to pyrolyses at 320°C. This type of lubricant has a temperature range of from minus 130 to plus 650°C.

Card 1/1







# WILSKI, I.


"The Ductube method of installing electric cables in concrete."

p. 46 (Wiadomosci Elektrotechniczne) Vol. 18, no. 2, Feb. 1958 Warsaw, Poland

SO: Monthly Index of East European Accessions (EEAI) LC. Vol. 7, no. 4, April 1958







MARCZEWSKI, K., mgr inz. arch.; WIISKI, Juliusz, inz. arch.;
WYPOREK, Bogdan, mgr inz. arch.

Organization and economics of the capital investment process. Przegl techn 35 no.31:4 2 Ag '64.

# WILSON, C.


SCIENCE

Periodicals MAGYAR FIZIKAI FOLYOIRAT Vol. 6, no. 4, 1958

WILSON, C. On the ionization of atmospheric air. Tr. from the English. p. hol.

Monthly List of East European Accessions (EEAI) LC. Vol. 8, No. 5, May 1959, Unclass.

erregies des regresses l'experientes les	是为65世纪的中国公司的公司的证据。	44 STATES
	$\chi$	
WILTHANSKA,		
	POL	
	U01,183.9 ; 511,163.24	
	Dasiássi A., Wilmaásku A. Intluence of Cleetrofyles on the Sarption of	
	Salphurous Acid by Polato Starch.	
	Woley elektrolitow na sorpely kwasu slarkuwego przez mączkę	
	ziemniaczany. Przemysł Chemiczny, Nu. 2, 1934, pp. 62-64, 3 figs.	
	5 tabs. The influence was investigated of electrolytes, NH.Cl, NaCl, KCl.	
	Lich, SrCl <sub>2</sub> , CaCl <sub>2</sub> , and MgCl <sub>2</sub> , on the sorption of sulphurous acid by po-	
	tatu starch in an anneous solution. It was established that the presence	
	nt there electrolytes raises the sarption, the increase being contingent to the increased concentration. No distinct differences of action of in-	
	dividual long in a 0.25 N electrolyte solution were observed. In a 0.5 N	
	solution, the progressively decreasing influence of ions can be represen-	
사용한 글로막으로 하는데 하는데 말라이라고 있는데 보다. 	ted us follows: $Cn + i > Mg + i > Sr + i > Li + i > K + i > Nn + i > NH_4 + i$ .	
	In higher electrolyte concentration the influence of magnesium lons is tronger than that of calcium ions. Glycerin and alcohol raise the ad-	
	corption rate; and the influence of these substances hereases with the	
	degree at concentration.	
	일하는 경우를 받는 것이 되었다. 	
	ENCOVER STANDARD STAN	
		CHECK THE RESERVE





HAGY, Z. WILTER, W.

Effect of isolated cerebral and isolated trunk hypoxia on minute volume. Acta physiol. hung. Suppl. no.6:55-56 1954.

1. III. Medizinische Klinik der Medizinischen Universitat, Budapest. (HEART minute volume, eff. of exper. cerebral or trunk hypoxia) (ANOXIA cerebral or trunk hypoxia, eff. on minute volume)

```
THE PROPERTY OF THE PROPERTY O
                                        GOMORI, Pal; SZABO, Gyorgy; KOVACS, Arisatid; NAGY, Moltan; TAKACS, Lajos;
WILTNER
                                                                                   WILTHER, Willibald
                                                                                                                 area de la companya d
                                                                                     The role of hypoxia and hypercapnia in the neuroregulation of the
                                                                                     renal circulation and diuresis. Magy Belorv. arch. 7 no.5:129-145
                                                                                       Oct. 54.
                                                                                        1. A Budapesti Orvostudomanyi Egyetem III. sz. Belklinikajanak
                                                                                         (igazgato: Gomori Pal egyetemi tanar) I. ss. Belklinikajanak
                                                                                         (igasgato: Ruswnyak Istvan egyetemi tanar) es Elettani Intezetenek
                                                                                          (Igazgato: Balint Peter eyetemi tanar) koslemenye.
                                                                                                                                     (CARBON DIOXIDE, in blood
                                                                                                                                                       hypercapnia, eff. on renal circ. & diuresis in dogs)
                                                                                                                                      (ANOXIA, exper.
                                                                                                                                                        eff. on renal circ. & diuresis in dogs)
                                                                                                                                       (KIDNEYS, blood supply
                                                                                                                                                         circ., eff. of exper. anoxia & mypercapnia in dog)
                                                                                                                                         (DIURESIS
                                                                                                                                                           eff. of exper. anoxia & hypercapnia in dogs)
```





WILTNER, Willibald, dr.

Oral therapy of diabetes mellitus with sulfamide derivatives.
Orv. hetil. 98 no.39:1081-1084 23 Sept 56.

1. A Budapesti Orvostudomanyi Egyetem III. sz. Belklinikajanak
(igazgato: Gomori, Pal; dr. egyet. tanar) kozlemenye.

(DIABETES MELLITUS, ther.

Nesulfanilyl-N'-butyl urea, indic. & mechanism of action (Hun))

(SULNOMANIDES, ther. use

Nesulfanilyl-N'-butyl urea, in diabetes mellitus, indic.
& mechanism of action (Hun))

(UREA, related cpds.

Nesulfanilyl-N'-butyl urea ther. in diabetes mellitus, indic. & mechanism of action (Hun))

UHAI, Leszlo, Dr.; NAGY, Zoltan, Dr.; SZINAY, Gyula, Dr.; WILTNER, Willibald, Dr.

Kidney examinations in scleroderma. Orv. hetil. 99 no.8-9:266-271 23 Feb

- 2 War 58.

1. A Budapesti Orvostudomany Egyetem III. sz. Belklinikajanak (igazgato:
Gomori Pal dr Egyet. tanar) kozlemenye.

(SCIENOMENMA, physiola.

kidneys, physiopathol. (Hun))

(KIDNETS, in various dis.

scleroderma, physiopathol. changes (Hun))

scleroderma, physiopathol. changes (Hun))

GOMORI, P.: KOYACH, A. G.B.; TAKAGS, L.; FOLDI, M.; SZABO, Gy.: NAGY, Z.;

WILTNEE, W.

Renal blood flow in arterial hypoxia. Acta med. hung. 16 no.1:
37-42 \*60.

1. 3rd Department of Medicine (Director: P. Gomori). Institute of Physiology (Director: P. Balint), and 1 st Department of medicine (Director: I. Rusznyak). University Medical School, Budapest.

(AMOXIA exper)
(KIDNEYS blood supply)

GOMORI, F.; KOVACH, A.G.B.; TAKACS, L.; FOLDI, M.; SZABO, Gy.; MAGY, Z.;


WILTHER, W.

The control of renal circulation in hypoxia. Acta med.hung. 16
no.1:43-60 '60.


1. 3rd Department of Medicine (Director: P.Gomori), Institute of
Physiology (Director: P.Balint), and 1 st Department of Medicine
(Director: I.Rusznyak), University Medical School, Budapest.


(AMUXIA exper)

(KIDNEYS blood supply)



Treatment of bronchial asthma in Kekesteto. Orv hetil 101 no.23: 814-816 5 Je '60.  1. Kekestetei Allami Gyogyintezet, II. Belosstaly. (ASTHMA ther.)			
814-816 5 Je '00.	WILLE	TER, Willibald, dr.	641 101 no.23:
Walter Allami Gyogyintezet, II. Belosstaly.		814-816 5 Je '60'-	
		Toboutetei Allami Gyogyintezet, II. Belosstaly.	






WILTOSINSKI, R.

SCIENCE

Periodicals: CHEMIK. Vol. 11, no. 7/8, July/Aug. 1958

WILTOSINSKI, R. Information on the mensuration equipment for the chemical industry planned and constructed by the Silesian Polytechnic School at Gleiwits. p. 3 of cover.

Monthly List of East European Accessions (EEAI) LC Vol. 8, No. 4, April 1959, Unclass.



WILUN, Z.

The geotectonic researches on the fundation of the Palace of Culture and Science in Warsaw. p. 237

Vol. 12, no. 7, July 1955 INZYNIERIA I BUDOWNICTWO Warszawa

Source: Monthly List of East European Accessions, (EEAL), IC, Vol. 5, no. 2
Feb. 1956

WILUN, Z.

WILUN, Z. Problems of soil mechanics as applied to surveying. p. 441.

Vol. 12, no. 12, Dec. 1956. PRZEGLAD GFODFZYJNY. Warszawa, Poland.


SOURCE: East European Accessions List (EEAL) Vol. 6, No. 4-April 1957

WILLIN, ZENON

Wyznaczanie dopuszczalnych obciazen gruntow

Warszawa, Poland, Arkady, 1958. 283 p.

Monthly List of East European Accessions (EEAI) LC, Vol. 8, no. 9, September 1959. Uncl.



WILUN, Zenon, prof. inz.; BOLENSKI, Marian, mgr inz.

Subsoil studies with the ITB-ZW percussive rotary soil trial tool. Inst tech budow inf no.12:40-42 '63.

1. Zaklad Mechaniki Gruntow i Fundamentowania, Instytut Techniki Budowlanej, Warszawa.


WILUN, Zenon, prof. dr inz.; KISIEL, Igor, prof. dr inz.; KWASNIEWSKI, Julian, mgr inz. HÜCKEL, Stanislaw, prof. dr inz.


Discussions on papers and communications. Rozpr hydrotechn no.12: 95-97 162.

1. Technical University, Warsaw (for Wilun). 2. Technical University, Wroclaw (for Kisiel). 3. Research Institute of Hydraulic Engineering, Polish Academy of Sciences, Gdansk (for Kwasniewski, Huckel).

"APPROVED FOR RELEASE: 03/20/2001 CIA-RDP86-00513R001961620001-5

WILUSZ, Zdzielaw	c/196h		1964
ECOLOGY	DECEASED		
	교통하는 경기를 가고 하고 있는 것이 되다. 대한민국의 교통 기계를 보고 있다. 대한민국의 대한민국의 기계를 보고 있다.	a di marana di Salaharan da Kababatan Balanda da Kababatan Balanda da Kababatan Balanda da Kababatan Balanda d Kababatan Balanda da Kababatan Balanda da Kababatan Balanda da Kababatan Balanda da Kababatan Balanda da Kabab	





WIMMER, M.

"Organizing the Work with Prefabricated Housing Constructions." p. 255 (Stavebni Prumysl, Vol. 3, no. 11, June 1953, Praha)

So: Monthly List of East European Accessions, Vol. 3, no. 2, Library of Congress, Feb. 1954, Uncl.

Construction of frame buildings. p. 62. POZEMNI STAVBY. (Ministerstvo stavebnictvi) Praha. Vol. 3, no. 2, Feb. 1955.

WIMMER, M.

SOURCE: East European Accessions List (EEAL), Library of Congress, Vol. 4, No. 12, December 1958.

Meeting of general secretaries of the Scientific Technical Societies in Czechoslovakia in 1962. Tech praca 14 no.4:247-249 Ap '62.  1. Vedecky tajemnik Ustredni rady Ceskoslovenske vedecko technicke spolecnosti.	WIMMER	, Miloslav
1. Vedecky tajemnik Ustredni rady Ceskoslovenske vedecko technicke spolecnosti.		Meeting of general secretaries of the Scientific Technical Societies in Czechoslovakia in 1962. Tech praca 14 no.4:247-249 Ap '62.
		l. Vedecky tajemnik Ustredni rady Ceskoslovenske vedecko technicke spolecnosti.

WIMMER, Miloslav, nositel Radu prace

International conference of scientists at the 4th Brno Fair. Tech praca 14 no.9:667-672 S '62.

1. Vedecky tajemnik Ustredni rady Ceskoslovenske vedeckotechnicke spolecnosti.

WIMMER,	Miloslav
	The Czechoslovak Scientific Technical Society is the main organizer of the development of science and technology. Tech praca 14 no.10:797-799 '62.
	1. Vedecky tajemnik, Ustredni rada Ceskoslovenske vedecko- technicke spolecnosti, Praha.
	도 있는데 그 이 그는 이 이 경기가 되었다. 그 그 사람이 하는 그 # 10 # 이 이 이 이 이 이 이 이 이 이 이 이 이 이 이 이 이 이

WIMER, Miroslav, nositel Radu prace

Looking forward to the 12th Congress of the Communist Party of Czechoslovakia. Elektrotechnik 17 no.12:333-334 D '62.

1. Vedecky tajemnik, Ceskoslovenska vedecko-technicka spolecnost, Praha.

# WIMMER, Miloslav Experiment in planning, management, and financing. Tech praca 16 no.2:91-93 F'64. Construction of a building by the method of raised floors. Ibid.:110-111 1. Vedecky tajemnik, Ustredni rada Ceskoslovenske vedeckotechnicke spolecnosti.

WIMMER, Miloslav

Quality has priority. Tech praca 16 no. 6:401-403 Je '64.

Reliability service. Ibid.:416-417

Effect of designer's education on the quality and technological standard of production and products. Ibid.:461-463

1. Scientific Secretary of the Central Council of the Czechoslovak Scientific and Technological Society, Prague.

MACAK, Jiri; WIMÆR, Petr; ZACHER, Jan

Analytic evaluation of the acid component of low-temperature tar from Most lignite, Sbor pal vod VSChT Vol. 5:251-279 '61 [publ; '62].

1. Katedra koksarenstvi a plynarenstvi, Vysoka skola chemickotechnologicka, Praha.

### 

1/30 8/058/62/000/009/061/069 A057/A101

9.3/20

AUTHORS:

Kazoń, Grażyna, Winawer, Irena

TITLE:

Activation of silver-magnesium secondary-electron emitters for

photomultipliers

PERIODICAL:

Referativnyy zhurnal, Fizika, no. 9, 1962, 49, abstract 9-3-97t

("Przegl. elektron.", 1962, v. 3, no. 1, 9 - 11, Polish)

TEXT: Emitters from an Ag-Mg (Ag 98%, Mg 2%) alloy with a coefficient of secondary emission of about 4, working at a 100 - 150 v voltage, are used in the two last stages of the photomultiplier developed in the Przemysl Institute of Electronics (Poland). The activation of the emitters is based on the formation of a magnesium oxide surface layer of 10<sup>-5</sup> - 10<sup>-4</sup> cm thickness. The maximum values for the coefficient of secondary emission is attained by oxidation of an Ag-Mg alloy in steam atmosphere. Curves are given which characterize the activation process and are taken under thoroughly controlled conditions in which the influence of side factors was excluded by preliminary washing and protracted heating of samples. A drawback of the silver-magnesium emitters is their low resis-

Card 1/2

Activation of silver-magnesium  S/058/62/000/009/061/069 A057/A101  tance to moisture, consequently they change strongly the emission properties during storage in air. Until the installation these emitters have to be stored in exiscators or evacuated ampullas. There are 4 references. [Przemyslowy. Inst. Elektroniki, PNR]  N. S.  [Abstracter's note: Complete translation]  Wydzial Lamp Obrazowyth, Zaklad Doswindczalny  PRZEmyslowego Instytutu Elektoniki  Card 2/2		
Activation of silver-magnesium  A057/A101  tance to moisture, consequently they change strongly the emission properties during storage in air. Until the installation these emitters have to be stored in exiccators or evacuated ampullas. There are 4 references. [Przemyslowy. Inst. Elektroniki, PNR]  N. S.  [Abstracter's note: Complete translation]  Wydzial Lamp Obersowyth, Zaklad Doswindczalny  PRZEmyslowego Instytutu Elektoniki		
Activation of silver-magnesium  A057/A101  tance to moisture, consequently they change strongly the emission properties during storage in air. Until the installation these emitters have to be stored in exiccators or evacuated ampullas. There are 4 references. [Przemyslowy. Inst. Elektroniki, PNR]  N. S.  [Abstracter's note: Complete translation]  Wydzial Lamp Obersowyth, Zaklad Doswindczalny  PRZEmyslowego Instytutu Elektoniki	en de la companya de La companya de la co	
tance to moisture, consequently they change strongly the emission properties during storage in air. Until the installation these emitters have to be stored in exsicoators or evacuated ampullas. There are 4 references. [Przemyslowy. Inst. Elektroniki, PNR]  N. S.  [Abstracter's note: Complete translation]  Wydzial Lamp Obrazowych, Zaklad Doswindczalny  PRZEmyslowego Instytutu Elektoniki		
during storage in air. Until the installation these emitters have to be stored in existing or evacuated ampullas. There are 4 references. [Przemyslowy. Inst. Elektroniki, PNR]  N. S.  [Abstracter's note: Complete translation]  Wydzial Lamp Obrazowych, Zaklad Doswiadczalny  PRZEmyslowego Instytutu Elektoniki	Activation of silver-magnesium AUD//AIUI	
[Abstracter's note: Complete translation]  Wydzial Lamp Obrazowych, Zaklad Doswiadczalny  PRZEmyslowego Instytutu Elektoniki	during storage in air. Until the installation these emitters have to be stored in exiccators or evacuated ampullas. There are 4 references. [Przemyslowy.	•
Wydział Lamp Obrazowych, Zakład DoswiaDczalny PRZEmysłowego Instytutu Elektoniki	N. S.	
Wydział Lamp Obrazowych, Zakład DoswiaDczalny PRZEmysłowego Instytutu Elektoniki		
PRZEMYSłowego Instytutu Elektoniki	[ADSTRUCTER S NOTE: Complete translation]	
PRZEMYSłowego Instytutu Elektoniki		3
PRZEMYSłowego Instytutu Elektoniki	Wydział LAMP OBRAZOWYCH Zakład DoswiADCZAMY	:
	PRZ Emyslaveas Institutu Elektoniki	
Card 2/2		
Card 2/2	그 이 아이 이 얼마요? 이 사이 되어 불편한 경험을 받아 되었다. 그리고 이 말음	
Card 2/2	그리는 사람들은 사람들은 항하면 되는 그 생각 바람은 작업이 가는데 모든 모든	
	Card 2/2	

# vimicki, J.

The behavior of petrographic constituents of Polish coals during crushing. p. 166.


KAKS, SMOLA, CAZ. Katowice, Poland Vol. 4, no. 4, July/Aug. 1959.

Monthly List of East European Accession. (EEAI) LC, Vol. 9, no. 1, Jan. 1960

Uncl.

"APPROVED FOR RELEASE: 03/20/2001 CIA-RDP86-00513R001961620001-5

ALLING ALLINGS	VED FOR REL	EASE: US/2U/	LECTRE	KDP80-005	 TOI/OOT DOI	77,085
				<del></del>	-37	
COUNTY CANADA		•			80590	
	jour. : REM	nine, No.22 1950 ila, W. and Win	tacklevion, '		p g. candidated com-	
OFFICE LANGE	toll :	Descropment of	f Specificar	TOUR AND THE	Polp ()	
		seglad Papiern  le authors have  aper rulp made	- ほんづきしつひをは 子は	311	しゅうがき されない	
A301	30 10 10	te mathors have apor rulp made acis of analyticased on the type pecifications (he busis for formations for a second formations for a second formation formation for a second formation for a second formation formation formation for a second formation for a second	cal data. The of parer	he classore the authors	mb =	
	20: 1/1		306	والإن المستويد والمستويد والمستويد والمستويد والمستويد المستويد والمستويد وا		



POLAND // Physical Chemistry. Radiochemistry.

Abs Jour: Ref Zhur-Khimiya, No 20, 1959, 70696.

: Zielinski, M.; Wincel, H. Author

Inst :Not given.

: Synthesis of Potassium Oxalate and Oxalic Acid Labeled with  ${\rm C}^{14}$ . Title

Orig Pub: Roczn. chem., 1958, 32, No 5, 1189-1190.

Abstract: In a previously proposed method of synthesizing K2C2O4 (I) labeled with Cl4 (Long E. A., J. Amer.

Chan. Soc., 1939, 61, 570), finely granulated sand (the adsorbent active agent) is replaced by gravel having a grain diameter of 5-7 mm. By the method of reverse isotope dilution, it is established that I, obtained by this method, contains no traces of carbonates. The labeled  ${\rm H_2C_2O_4}$  (II) was obtained by means of isotope ex-

Card 1/2

16

CIA-RDP86-00513R001961620001-5" **APPROVED FOR RELEASE: 03/20/2001** 

POLAND / Physical Chemistry. Radiochemistry. Isotopes. B

Abs Jour: Ref Zhur-Khimiya, No 20, 1959, 70696.

Abstract: change between inactive II and active I or by

means of treating I with inorganic acid, followed by an ether extraction of II. B. Kaplan.

monthly hist of East European (eccesions). (EEAI) LC. Vol. 8, No. 7, Jul. 59 Unel.

Card 2/2

ZLOTOWSKI, Ignacy; WINCEL, Henryk

Mass spectrometric studies of chemical processes occurring in a selfquenching Geiger-Muller counter filled with long-chain saturated hydrocarbons. Mukleonika 5 no.5:261-279 '60.

1. Warsaw University, Warszawa, Department of Muclear Chemistry

ZLCTCWSKI, Ignacy; WINCEL, Henryk

Mass spectrometric studies of electron impact fragmentation of ethyl halides. Nukleonika 6 no. 6:409-414. '61

1. Warsaw University, Warszawa, Department of Nuclear Chemistry.

ZLOTOWSKI, Ignacy; POLACZEK, Andrzej; WINCEL, Henryk

A study of the icdine exchange between KAII, and alkyl halides by the method of isotopic tracers. Nukleonika 6 no. 6:415-422. '61

1. Universytet Warszawski, Warszawa, Katedra Chemii Jadrowej.

33736

11.3950 11.1210 5.4600

P/046/62/007/001/002/006 D216/D304

AUTHOR:

Wincel, Henryk

TITLE:

Mass spectra and primary radiation processes of certain

organic substances in the gas phase

PERIODICAL:

Nukleonika, v. 7, no. 1, 1962, 25-38

TEXT: Using the quasi-equilibrium theory of H. Eyring and his associates the author calculated the distribution of unimolecular decomposition products of irradiated methane and ethane, and G-values for the methane products. The time interval used in the calculations was the average time between gas molecule collisions at atmospheric pressure 10<sup>-10</sup> sec. The activation energies of the possible unimolecular reactions were estimated from appearance potentials or thermochemical calculations as given by F.N. Field and J.L. Franklin and M. Krauss, A.L. Wahrhaftig and H. Eyring. The mass spectrum for methane was calculated using the expression Eq.(3)

 $P(E) = e^{-\frac{27}{2}\left(\frac{E}{E^2}\right)}$  with P(E)=0 from 5-10 eV, E being the average excitation energy. For ethane, P(E)=1 for  $0 \le E \le 6.6$  eV and E = 0 for E = 0 fo

33736 P/046/62/007/001/002/006 D216/D304

Mass spectra and primary ...

Card 2/4

E 7 6.6 eV. The calculated relative abundances of unimolecular decomposition products in both cases differ considerably from experimental data, indicating that bimolecular ion-molecule reactions complicate the distribution obtained from experimental mass spectra. The author now assumes that primary products undergo bimolecular reactions in collisions with molecules of the parent substance, and that these reactions occur at every ion-molecule collision. It is shown that, taking this into account, the ion ratios  $\text{CH}_5^+/\text{C}_2^{\text{H}_5^+}$ ,  $\text{CH}_5^+/\text{C}_2^{\text{H}_4^+}$ , and  $\text{C}_2^{\text{H}_5^+}/\text{C}_2^{\text{H}_4^+}$  calculated for methane are in satisfactory agreement with experimental data. All these ions have high yields, and greater deviations were found for the low yield products. Considering radical and atom reactions, only the CH<sup>+</sup> ion is assumed to decompose on neutralization. It is shown that in addition to recombination and disproportionation reactions, in which free radicals and hydrogen atoms take part, the reaction of hydrogen with methane must be considered; the reaction of free radicals with the methane may, however, be neglected. From all the possible reactions, the results were calculated, taking W=27,3 eV/ion pair. The statistical method used by J.H.

33736 P/046/62/007/001/002/006 D216/D304

Mass spectra and primary ...

Futrell and C.D. Wagnet and kinetic data of C.A. Heller, S. Rabinovitch et al. J.A. Kerr and A.F. Trotman-Dickenson and K.J. Ivin and E.W. R. Steacie were employed in the calculations. Comparison with the experimental results is difficult due to the differences of methods and dependence of the degree of conversion of the parent substance. The results of G.J. Mains and A.S. Newton were used and the relative G-values were compared with the relative experimental yields of the appropriate products. Good agreement is obtained, supporting the method of radiolysis proposed by the author, and indicating the possible value of the quasi-equilibrium theory. The author thanks Professors I. Zlotowski, S. Minc, and Doctor Kecki. There are 2 figures, 7 tables and 44 references: 4 Soviet-bloc and 40 non-Soviet-bloc. The 4 most recent references to the English-language publications read as follows: J.H.Futrell. J.Phys.Chem., 64,1634,1960; A.Korf, E.M. Eyring, A.L. Wahrhaftig, H. Eyring: J. Chem. Phys. 32, 149, 1960; E.M. Eyring, A.L. Wahrhaftig: J. Chem. Phys., 34, 23, 1961; G.J. Mains, A.S. Newton, J. Phys. Chem., 65, 212, 1961.

Card 3/4

### "APPROVED FOR RELEASE: 03/20/2001

33736

P/046/62/007/001/002/006 D216/D304

Mass spectra and primary ...

Institute of Nuclear Research, Warsaw, Department of

Radiation Chemistry

SUBMITTED:

ASSOCIATION:

November, 1961

Card 4/4

CIA-RDP86-00513R001961620001-5" APPROVED FOR RELEASE: 03/20/2001

WINCEL, Henryk

Mass spectra and primary radiation processes of certain organic substances in gas phase. Nukleonika 7 no.1:25-38 '62.

1. Institute of Nuclear Research, Warszawa, Department of Radiation Chemistry

Determination of radiation yield of hydrogen by the gas chromatography method. Mukleonika 7 no.3:169-174 162.
1. Polish Academy of Sciences, Institute of Nuclear Research, Department of Radiation Chemistry, Warsaw.

AUTHORS: Zagórski, Zbigniew, P., Kecki, Zbigniew, Wincel, Henryk

TITLE: Determination of radiation yields in the two-phase system of O2(gas)-H2O, NaOH, Na2SO3(liquid). I.

PERIODICAL: Referativnyy zhurnal. Khimiya, no. 1, 1963, 88, abstract 1B611 (Rept. Inst. badan jedrow. PAN, 1961, N 290/ChR, 7 pp., ill. [Eng.; summaries in Pol. and Russ.])

TEXT: The yields of G(-SO3), G(-O2), and G(H2) were determined in the two-phase system: gaseous O2 - 0.066 M Na2SO3 solution - 1 M MaOH under

TEXT: The yields of  $G(-SO_3^-)$ ,  $G(-O_2)$ , and  $G(H_2)$  were determined in the two-phase system: gaseous  $O_2$  - 0.066 M Na<sub>2</sub>SO<sub>3</sub> solution - 1 M HaOH under the effect of  $CO_2^{60}$   $\gamma$ -irradiation.  $G(H_2)$  = 0.44  $\pm$  0.02,  $G(-SO_3^{2-})$  and  $G(-O_2)$  decrease when the dose increases,  $G(-SO_3^{2-})/G(-O_2) \approx 2$ . [Abstracter's note: Complete translation.]

Card 1/1

ACCESSION NR: AP3003624

P/0046/63/008/002/0117/0127

AUTHOR: Kecki, Zbigniew; Wincel, Henryk

TITLE: Primary processes of radiation chemistry as studied by mass spectrometry. I. Fragmentation of tetrahydronaphthalene on electron impact

SOURCE: Nukleonika, v. 8, no. 2, 1963, 117-127

TOPIC TAGS: tetrahydronaphthalene electron fragmentation, electron impact fragmentation, tetrahydronaphthalene, tetrahydronaphthalene fragmentation, radiolysis metastable ion, radiolysis primary process, molecular fragmentation

ABSTRACT: A mass-spectrometric investigation of the fragmentation of tetrahydronaphthalene under electron impact has been carried out. "Pure" grade domestic tetrahydronaphthalene was dried and purified in a column of 15 theoretical plates. The fraction used had a boiling point of 207C at 754 mm Hg. Gas chromatrography and mass spectrometry analysis showed the presence of about 0.5% decahydronaphthalene. "Analytical" grade domestic toluene was used without extra purification, as gas chromatography revealed no impurities. Both substances Cord 1/2

	<u> </u>	· · · · · · · · · · · · · · · · · · ·
CCESSION NR: AP3003624		
eing introduced into the session of	al freezing and pumping cycles in a ion source of the mass spectroled by a 60° Soviet mass spe	meter (des- s were observed, ragmentation Large quan-
erved had a cyclic formons produced from other add 2 tables.	a and decomposed in the same way organic substances. Orig. art	. has: 3 figures
SSOCIATION: Institute hemistry, Warsaw, Polar UBMITTED: 180ct62	DATE ACQ: 19Ju163 ENC	L: 00
JB CODE: CH.PH	NO REP SOV: 001 OTH	BR: 034
rd 2/2		

WINCEL, Henryk; KECKI, Zbigniew

Primary processes in radiation chemistry as studied by mass spectrometry. Pt. 2. Mukleonika 8 no.4:215-223 363.

1. Institute of Nuclear Research, Department of Radiation Chemistry, Warsaw 9.

WINCEL, Henryk; KECKI, Zbigniew

Primary processes in radiation chemistry as studied by mass spectrometry. Pt.3. Nukleonika 8 no.8:529-543 '63.

1. Department of Radiation Chemistry, Institute of Nuclear Research, Warsaw.

L 20588\_66 EMP(j)/FMA(h) RM

ACC NR: AP6012010 SOURCE CODE: PO/0046/65/010/09-/0567/0573

AUTHOR: Wincel. Henryk—Vintsel', G.; Kecki, Zbigniew—Kentskly, Z. 

ORG: Department of Radiation Chemistry, Institute of Nuclear Research, Warsaw-Zeran

TITIE: Primary processes in radiation chemistry as studied by mass spectrometry. IV.

Ion-molecule reactions in tetrahydronaphthaleney

SOURCE: Nukleonika, v. 10, no. 9-10, 1965, 567-573

TOPIC TAGS: radiation chemistry, mass spectrometer

ABSTRACT: Ion-molecule reactions in which neutral end ionized molecules of tetrahydronaphthalene take part were investigated with a mass spectrometer. Relative cross-section values of individual reactions were determined and the reactions occurring in the radiolysis process of tetrahydronaphthalene in the liquid phase are discussed.

Orig. art. has: 2 figures, 7 formulas, and 3 tables. NA

SUB CODE: 07 / SUEM DATE: 19Jun65 / ORIG REF: 005

# SKEELIN		
	L 33005-66 EWP(j) .RM	
	ACC NR: AP6024167 SOURCE CODE: PO/0046/65/010/012/0753/0764	
	AUTHOR: Wincel, HVintsel', G.; Kecki, Zbigniew-Kentski, Z.	
	ORG: Department of Radiation Chemistry, Institute of Nuclear Research, Warsaw-Zeran	
	TITLE: Primary processes in radiation chemistry as studied by mass spectrometry.  V. Excited and superexcited states in the radiolysis of tetrahydronaphthalene	
	SOURCE: Nukleonika, v. 10, no. 12, 1965, 753-764	
	TOPIC TAGS: radiation chemistry, mass spectrometry, excited state, chemical decomposition, recombination reaction	
	ABSTRACT: The formation of ions and excited molecules in the initial stage of radic-lysis of tetrahydronaphthalene was analyzed. Their yields were determined to be:  G(C10H2=4.0 and G(C10H2)=4.3. The possibility of formation of excited and superexcited molecules in the recombination process and the transformations of these molecules are discussed. The yields of products of the transformations were evaluated. The authors thank Professor, Doctor Stefan Minc for his guidance and interest in this work. Orig. art. has: 8 formulas and 2 tables. Orig. art. in Eng.	
	SUB CODE: 07 / SUBM DATE: 140ct65 / ORIG REF: 008 / SOV REF: 005	
	Card 1/1 Ala 1767	